

WO 92/05194(1)
A61K39/10-C07K15/00F14A24-
[M07K201:02]-

PROPERTY ORGANIZATION
al Bureau



DER THE PATENT COOPERATION TREATY (PCT)

-2- * -C07K15/00F14A24

14/235

C07K 13/00, A61K 39/10

A1

(11) International Publication Number: WO 92/05194

(43) International Publication Date: 2 April 1992 (02.04.92)

(21) International Application Number: PCT/NL91/00185

(22) International Filing Date: 25 September 1991 (25.09.91)

(30) Priority data:
9002092 25 September 1990 (25.09.90) NL

(71) Applicant (for all designated States except US): DE STAAT
DER NEDERLANDEN VERTEGENWOORDIGD
DOOR DE MINISTER VAN WELZIJN, VOLKSGE-
ZONDHEID EN CULTUUR [NL/NL]; P.O. Box 5406,
NL-2280 HK Rijswijk (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HAMSTRA, Hendrik-
Jan [NL/NL]; Rosmolen 36, NL-3961 LJ Wijk Bij Duur-
stede (NL). POOLMAN, Jan, Teunis [NL/NL]; Lee-
teinde 8, NL-1151 AK Broek in Waterland (NL).

(74) Agents: DE BRUIJN, Leendert, C. et al.; Nederlandsch
Octrooibureau, Scheveningseweg 82, P.O. Box 29720,
NL-2502 LS The Hague (NL).

(81) Designated States: AT (European patent), BE (European
patent), CA, CH (European patent), DE (European pa-
tent), DK (European patent), ES (European patent), FR
(European patent), GB (European patent), GR (Euro-
pean patent), IT (European patent), LU (European pa-
tent), NL (European patent), SE (European patent), US.

Published

With international search report.

*Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: VACCINE SUITABLE FOR COMBATting BORDETELLA PERTUSSIS

(57) Abstract

The invention relates to vaccines, which are suitable for combatting *Bordetella pertussis*, the causative organism of whooping cough. More in particular, the vaccines comprise, as active component, one or more outer membrane proteins (OMPs) derived from *Bordetella pertussis* or from genetically manipulated microorganisms producing said OMPs. Preferably the OMPs having a molecular weight of about 32 kDa and 92 kDa - either separate or as a combination of both - are applied as the active component. According to an embodiment of the invention the OMPs in question are present in an outer membrane vesicle (OMV) formulation or in an artificial vesicle formulation like a protein-detergent formulation.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU ⁺	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE ⁺	Germany	MC	Monaco	US	United States of America
DK	Denmark				

⁺ Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

Vaccine suitable for combating Bordetella pertussis.

The invention primarily relates to a vaccine which is suitable for combating B.pertussis in humans.

5 Prior art

As reported in The Lancet, June 2, 1990, pp. 1326-1329, Bordetella pertussis is the causative organism of whooping cough. More particularly, B.pertussis is a Gram-negative bacterium which adheres to the cilia of epithelial cells in the human respiratory tract, where the bacterium reproduces and gives off toxic substances. At the local level, the infection causes destruction of the epithelium of the respiratory tract and kills the cells provided with cilia, which gives rise to, for example, impeded breathing, paroxysmal coughing, apnoea and encephalopathy, which may or may not be accompanied by fever. The infection can occur in humans at any age but occurs mainly in infants and young children.

Although B.pertussis is sensitive to a variety of antibiotics, treatment with such antibiotics is not effective since, before the disease is diagnosed, the B.pertussis bacteria have often already attacked the respiratory tract and have given off the toxins which are responsible for the serious consequences brought about by the disease. Therefore, preventive protection against B.pertussis is a matter of crucial importance.

In the light of the above, vaccines have been developed, the "whole-cell" vaccine of B.pertussis having been used as first vaccine. A vaccine of this type consists of whole B.pertussis cells, which have been killed, for example by the action of heat or a formalin treatment. This type of vaccine is effective in preventing whooping cough. Nevertheless, the use of this type of vaccine is highly controversial because of the occurrence of local and systemic side effects. Some effects are mild reactions, which also occur with other vaccines, but nevertheless occur more frequently in the case of B.pertussis vaccines. Examples of such effects are redness, pain, induration and fever. The B.pertussis "whole-cell" vaccine also causes other reactions specific for this type of vaccine, such as persistent crying and convulsions. Moreover, severe side effects such as brain damage and even death also arise with this type of

vaccine.

In the light of the above-described disadvantages associated with B.pertussis "whole-cell" vaccines, which vaccines contain a variety of substances such as, for example, proteins, nucleic acids, peptidoglucans, lipids and lipopolysaccharides, a broad-based study has been started to develop acellular vaccines, with which, in the optimum case, only those antigens which generate adequate immunity and themselves display no toxicity should be allowed to be present.

The generally recognised intracerebral (i.c.) mouse protection test [Medical Research Council. 1956, Vaccination against whooping cough. Relation between protection tests in children and results of laboratory tests. Brit.Med.J. 2: pp. 454-462] is chosen as a criterion for the effectiveness of such acellular vaccines.

As can be seen from the abovementioned article in The Lancet, vaccines containing various antigens have been tested. One of the examples is detoxified pertussis toxin (PT toxoid) which apparently - according to the i.c. mouse protection test - is less effective than the "whole-cell" vaccine. Only when one or more other antigens, such as, for example, the outer membrane protein (OMP) having a molecular weight of 69kDa (69K OMP) are added is a potency achieved which to some extent approaches that of the "whole-cell" vaccine. One example of such acellular vaccines is, inter alia, a mixture of filamentous haemagglutinin (FHA) and pertussis toxin (PT), which have been obtained together from culture supernatants and detoxified with the aid of formaldehyde. However, there is always the risk with the use of such detoxified PT-containing vaccines that a reversion to toxicity takes place during the clinical treatment. For this reason, genetically deactivated forms of pertussis toxin have been developed. In this case, the B.pertussis toxin gene has been modified by means of deletion, insertion or substitution of codons coding for specific amino acids in such a way that the PT then has hardly any toxicity but has retained the immunity-generating action [Pizza, M.G. et al., Mutants of pertussis toxin suitable for vaccine development. Science 246. (1989), pp. 497-500].

INVENTION

In view of the problems described above, the Applicant has carried out a study to find effective, non-toxic acellular B.pertussis vaccines which - after administration - provide an effective protection against B.pertussis in humans.

Vaccines suitable for combating B.pertussis have been found, which vaccines comprise, as active components, at least one or more outer membrane proteins (OMPs) derived from B.pertussis. Preferably these OMPs are present in an outer membrane vesicle (OMV) formulation or an artificial vesicle formulation. Examples of artificial vesicle formulations are, for example, a liposomal, proteosomal and detergent micelles.

Vaccines according to the invention advantageously comprise OMPs respectively OMVs or artificial vesicles, containing OMPs, which are derived from B.pertussis strains in the virulent stage i.e. OMPs having a molecular weight of about 92kDa and 32kDa (92K OMP and 32K OMP). In the i.c. mouse protection test, vaccines of this type give values which are well above the internationally specified minimum protection value of 4. Further studies carried out by the Applicant have shown, surprisingly, that a relationship exists between the ratio between the amounts of OMPs having a molecular weight of 92kDa and 32kDa (92K OMP and 32K OMP) and the OMP having a molecular weight of 38kDa (38K OMP), on the one hand, and the efficacy or protection value (according to the i.c. mouse protection test), on the other hand. On the basis of these findings, the 92K OMP/38K OMP weight ratio must, according to the Applicant, be at least greater than 0.25 and advantageously greater than or equal to 0.4, for example 0.5 or even higher. The relationship concerned is shown as a graph in Fig. 1. OMPs as well as OMVs having an optimum OMP-ratio can be obtained by means of genetic engineering. OMPs as well as OMVs having optimum ratios of 92kDa and 32kDa OMPs relative to the 38kDa OMP, which is indispensable for growth, can be prepared by this generally known technique.

Furthermore, the invention advantageously relates to artificial vesicles in which, preferably, at least the 32kDa OMP and/or the 92 kDa OMP is present. An OMP of this type can be brought into the form of so-called "mixed" protein-detergent micelles with

the aid of a detergent such as Zwittergent 3-14, and also other detergents like Triton X-100 and octylglucoside. According to the i.c. mouse protection test, artificial vesicles of this type have valuable activities, as is indicated below in Table 3.

5 Also other ingredients which are suitable as vaccine ingredients can be added to the vaccines in question. Examples of such ingredients are for example adjuvants like $AlPO_4$.

With a view to the presence of lipopolysaccharides in OMVs, which (can be) are active as endotoxins, the invention also relates to vaccines which comprise OMVs which have been treated with an effective amount of desoxycholate in order to lower the endotoxin content.

10 Better definable vaccines according to the invention can be obtained with the aid of fimbriae and/or FHA B.pertussis strains. Such strains can be selected in a simple and generally known manner.

The invention also relates to vaccines whose efficacy has been increased by the addition of additional active components such as PT toxoid and/or FHA. Although the PT toxoid can be derived from PT detoxified using formaldehyde or glutaraldehyde, the PT toxoid preferably used is that which has been extracted from B.pertussis strains in which the PT gene has been modified in such a way that it yields a non-toxic PT product of adequate immunological activity.

20 The vaccines according to the invention are used for the vaccination of humans, in particular children less than 2 years old. The dose to be injected can be about 5-25 μg OMV.

Finally, the invention relates to the above-defined OMPs and OMVs, derived from B.pertussis strains, which can be used for the preparation of the vaccines according to the invention.

30

Detailed description of the invention

Many B.pertussis antigens, such as detoxified PT, filamentous haemagglutinin (FHA) fimbriae, virulent stage OMPs and avirulent stage OMPs were reviewed in the study carried out by the Applicant. All of these antigens were tested using the intracerebral (i.c.) mouse protection test to determine their activity. This is because it has been found that this internationally known i.c. mouse

35

protection test is correlated to the field protection.

5 The abovementioned antigens, which differ from PT toxoid, were obtained from a PT-negative mutant of B.pertussis, specifically the BP-TOX6 mutant, as described in Relman, D.A. et al., Filamentous hemagglutinin of Bordetella pertussis: nucleotide sequence and crucial role in adherence. Proc.Natl.Acad.Sci 86 (1989), pp. 2637-2641. This BP-TOX6 mutant lacks the pertussis toxin operon. More particularly, the following antigens were obtained from this mutant: purified FHA, fimbriae, OMVs and 92K OMP, 69K OMP, 38K OMP and 32K
10 OMP.

The results of the B.pertussis antigens tested and of a "whole-cell" B.pertussis vaccine (WCV) are given in Table 1 below.

TABLE 1

	KH 85/1 (whole-cell vaccine)	6.0
	Pertussis toxoid (formaldehyde)	12.8
	Pertussis toxoid (9K/129 G mutant)	14.9
5	FHA (BP-TOX6)	11.8
	OMV (BP-TOX6)	6.3
	OMV/DOC (BP-TOX6)	4.3
	OMV (BP-TOX6, C mode)	1.3
	OMV (BP 509)	8.1
10	OMV (BP 509, C mode)	2.7
	OMV (BP 509, fim ⁻ mutant)	9.8
	OMV (BP 509, FHA ⁻ mutant)	6.2
	OMV (BP 134)	9.5
	OMV (BP 134, C mode)	3.1
15	OMV (BP W28)	3.1
	OMV (BP W28, 69K ⁻ mutant)	5.0

fimbriae 2+3 ND

69K OMP ND

20 92K OMP ND

38K OMP ND

32K OMP ND

ND = not detectable

25 C mode: avirulent stage

fim⁻ mutant: mutant, which has no fimbriae

FHA⁻ mutant: mutant which has no FHA

DOC: subjected to desoxycholate treatment

30 As can be seen from Table 1 above, a number of antigens have a protection value of ≥ 4 . These include PT toxoid (either treated with formaldehyde or the mutated non-toxic PT), FHA and OMV. The purified fimbriae and OMPs having a molecular weight of 92kDa (92K OMP), 69kDa (69K OMP), 38kDa (38K OMP) and 32kDa (32K OMP),
 35 however, had no measurable protection value.

Tests on various preparations, analysed using SDS-PAGE and electroblotting after colouring with monoclonal antibodies (anti-PT,

anti-FHA, anti-fimbriae, anti-92K OMP, anti-69K OMP, anti-38K OMP and anti-32K OMP) for the specific proteins or Coomassie Blue for the total protein, showed that

- the WCV preparation essentially contains outer membrane proteins,
- 5 - OMV in the avirulent stage contains 38K OMP, 33K OMP and 18K OMP, and
- OMV in the virulent stage contains 92K OMP, 32K OMP and 30K OMP.

It can be seen from the appended Fig. 2 that the avirulent stage OMPs (38, 33 and 18kDa) can be differentiated from the
10 virulent stage OMPs (92, 32 and 30kDa) on the basis of trypsin sensitivity and Triton X-100/MgCl₂ extractability.

The following is pointed up with regard to the endotoxin content present in the preparations tested. The endotoxin content in the FHA preparation was considered high by the Applicant. The WCV
15 and OMV vaccines have comparable endotoxin contents. It proved possible to lower this endotoxin content of the OMV preparations by a factor of 10⁴ using the desoxycholate (DOC) treatment described in more detail below. This DOC treatment gave rise to a reduction in vesicle size of 40-180 nm (OMV) to 10-25 nm (OMV-DOC).

20 The results of serological analysis of antibodies generated in mice, using a single dose immunisation with WCV, OMV, PT_x, FHA, fimbriae, 92K OMP, 38K OMP and 32K OMP, are shown in Table 2 below.

TABLE 2

Antibody response in mice immunised with WCV, OMV, fimbriae, FHA and PT_x

		<u>ELISA antigen</u>						
	<u>Vaccine</u>	WC*	OMV	LPS	fim 2	fim 3	PT	FHA
5	WCV 12.5	1.721	2.221	0.093	0.085	0.038	0.030	0.046
	PT _x (form)	0.750	0.871	0.059	0.110	0.097	2.310	0.479
	PT _x (mut.)	0.305	0.282	0.048	0.080	0.072	2.996	0.285
	FHA	0.498	1.031	0.026	0.018	0.066	0.043	2.443
	OMV	2.616	3.270	0.648	0.633	0.185	0.038	0.699
10	fim 2+3	2.508	3.169	0.037	1.826	1.075	0.217	0.063
	92K	2.035	1.986	0.056	0.188	0.102	0.178	0.069
	38K	1.484	2.399	0.024	0.146	0.057	0.095	0.117

* "Whole-cell" preparation for ELISA testing was prepared from shaking-bottle cultures.

It can be deduced from the above Table 2 that the prepared WCV vaccine (in the virulent stage) yields no detectable antibody formation against PT, FHA and fimbriae. PT_x, FHA and fimbriae induce antibodies, as detected by WC and OMV. The WC and OMV preparations, used as coating antigen, contain some FHA and fimbriae, while FHA and fimbriae preparations contain certain OMP.

OMV vaccines induce some antibodies against fimbriae and FHA. In order to prevent this effect, the Applicant used fimbriae⁻ and FHA⁻ variants of the 509 strain, obtained by means of colony blotting using monoclonal antibodies, in its study. Variants of this type are illustrated in Fig. 3. OMV vaccines, prepared from such variants, had protection values similar to those of OMV vaccines prepared from the wild type 509 strain.

OMV vaccines prepared from other tested B.pertussis strains in the virulent stage, such as the PT⁻ strain BP-TOX6, strain 134 and the 69kDa⁻ mutant of the W28 strain, have an adequate i.c. mouse protection value. When the B.pertussis strains BP-TOX6, 509 and 134 were in the avirulent stage, the i.c. mouse protection value of the OMV vaccines decreased sharply. In this context it is emphasised that the difference between the OMVs in the virulent and the avirulent stage lies in the presence,

discussed above, of the 92K OMP, 32K OMP and 30K OMP in the virulent stage. The greater the 92K OMP and 32K OMP content relative to the 38K OMP content, the greater is the i.c. mouse protection value of the OMV vaccines (see Table 1 and Fig. 1). The preference for a high 92K OMP/38K OMP ratio in OMV vaccines is clearly apparent from Fig. 1, which shows the critical value of 4 with regard to the protection potency. Above aspect may also be expressed in the 32K OMP/38K OMP ratio as the 92K OMP/32K OMP ratio is constant in non-manipulated B.pertussis strains (see Figure 3). In Table 3 below, the results of experiments i.e. the i.c. mouse protection test are reported which are performed with purified 92K OMP, 38K OMP and 32K OMP in a Zwittergent-3,14 solution in the presence and absence respectively of pertussis toxin (PT). From the results of these experiments with these artificial vesicles it can be deduced that the purified 32K OMP and 92K OMP in a correct formulation provide a sufficient activity against a B.pertussis challenge.

TABLE 3

i.c. potency of purified OMPs

Antigen	PT	Surviving mice (10)
92K + Zw* 3-14	-	2/2/2/4
92K + Zw 3-14	+	8/8/5/1
32K + Zw 3-14	-	8/5/1/2
32K + Zw 3-14	+	10/8/7/4
38K + Zw 3-14	-	1/3/4/6
38K + Zw 3-14	+	4/8/3/1

* Zw = Zwittergent

In the above table, column 1 shows the antigen used, column 2 the presence (+) of PT or absence (-) of PT and column 3 the survival rate of the mice (10 per test).

The amino acid sequences of a few OMPs, which were determined with the aid of a gas phase sequencer, are given below.

- (a) 92kDa OMP
AAVTAAQRIDGGAAFLGDVAIAT(T)K(A)(S)(E)
(A)
- (b) 32kDa OMP
5 ALSKRMGELRLTPVAGGV(W)(G)(R)AF(G)
(V)
- (c) 33kDa OMP
10 ALSKRMGELRLTPVAGGVWGRAFV (G)(Y)(Q)
(G)(R)(R)(V)
(N) (L)
- (d) 30kDa OMP
ALDKRLGELRL(N)A DAG (G)---
(P)
- (e) 38kDa OMP
15 (E)TSVTLYGIIDTGI(G)YNDV(D)FKV(K)GANA-(D)
(T)

In view of the above it is considered desirable to produce the 92 kDa and 32 kDa OMPs preferably by B. pertussis or optionally other
20 microorganisms. This production can be realized by cloning the relevant structural genes and expressing these genes at an increased level; this last-mentioned production method can be considered standard technology. An example of the above standard technology is elucidated in Figure 4. More in particular Fig. 4
25 illustrates the well-known plasmid pUC18 having a 3.8 kb insert comprising the 32 kDa OMP gene. This structural gene is obtainable from the gene bank (Mooi et al, 1987, Microbial. Pathogenesis, 2, pp. 473-484) with the help of a nucleotide sequence derived from the N-terminus of the 32 kDa OMP:
30 5' ACCCCGGTCGCCGGCGGCGTGTGGGGCCGCGCCTTC 3'

MATERIALS AND METHODS

Strains and growth conditions

35 The Bordetella pertussis strains 134 (atypical LPS, fim 3) and 509 (typical LPS, fim 2) are the two strains which are used to prepare the Dutch "whole-cell vaccine" (WCV) (P.A. van Hemert, Specific properties of acid precipitated pertussis vaccines. Progr.Immunobiol. Standard 3, (1969), pp. 297-301).

The Tohama strain was supplied by Ch. R. Manclark (FDA, Bethesda, USA).

5 The BP-TOX6 strain was supplied by R. Rappuoli (Sclavo Research Center) and has already been described in Relman, C.A. et al., Filamentous hemagglutinin of Bordetella pertussis: nucleotide sequence and crucial role in adherence, Proc.Natl.Acad.Sci. 86 (1989), pp. 2637-2641.

10 The W28 strain and W28 (69K⁻) strain were supplied by G. Doughan (Wellcome Ltd., England). In this case the 18323 strain was used as the control strain (see Relman et al., Proc.Natl.Acad.Sci. 86 (1989), p. 2637, Sato et al., Infect.Immun. 41 (1983), p. 313 and Kendrick et al., Am.J.Publ.Health 37 (1947), p. 803).

15 Mutants of strain 509 were obtained with the aid of a colony blotting using monoclonal antibodies. After culturing for three days on Bordet-Gengou agar plates (Difco) the colonies were blotted on nitrocellulose filters and examined for the presence of fimbriae, FHA and 92kDa OMP. The fimbriae⁻ and FHA⁻ mutants were easily detected in this way.

20 Small-scale growth of B.pertussis in fluid was carried out in shaking bottles containing Verwey medium: 14 g/l casamino acid, 0.2 g/l KCl, 1 g/l starch, 0.5 g/l KH₂PO₄, 0.1 g/l MgCl₂.H₂O, 0.02 g/l nicotinic acid and 0.01 g/l glutathion. 0.5 g/l nicotinic acid was added for growth in the avirulent C mode.

25 Large-scale fermentation in fluid was carried out in 40 l, 140 l or 300 l Bilthoven units in B2 medium with pH and PO₂ control (P.A. van Hemert, Specific properties of acid precipitated pertussis vaccines. Progr.Immunobiol.Standard 3 (1969), pp. 297-301).

30 The cells were deactivated by heating at 56°C for 10 min., centrifuged at 3000 g and resuspended in a physiological saline solution.

35 Opacity units were detected with the aid of a reference standard. The protein determination was carried out using the BCA protein analysis reagent (Pierce) with BSA as standard.

Intracerebral (i.c.) mouse potency test

The i.c. test was carried out using the method of Kendrick et al. (Kendrick, P.L. et al., Mouse protection test in the study of pertussis vaccine, Am.J.Publ.Hlth. 37, (1947), pp. 803 ff). Male and female NIH-bred mice (10-14 g) were used.

Each vaccine was tested in four fivefold-diluted solutions (20 mice/solution) by i.p. immunisation (0.5 ml, containing 1 mg of AlPO_4). WCV was tested with 1 OU, 0.2, 0.04 and 0.008 OU.

Since 1 OU contains approximately 15 μg of protein, the final calculations were made for purified antigens on a comparable protein basis.

OMV, PT_x and FHA were tested in 5, 1, 0.2 and 0.04 μg doses and purified antigens in 25, 5, 1 and 0.2 μg doses on the basis of trial experiments. A WCV reference of standard potency is part of each experiment.

14-16 days after immunisation, the mice were injected intracerebrally with 10 μl of a suspension of B.pertussis 18323, containing 15×10^3 bacteria. This suspension was prepared from a lyophilised culture having a LD_{50} of $< 10^3$ bacteria. This culture was cultured for 4 days on Bordet-Gengou plates and then cultured for a further 24 hours on Bordet-Gengou and suspended in 1 part by weight of casamino acid in a physiological saline solution to obtain a concentration of 15×10^5 bacteria/ml.

Twelve mice were each injected with 15×10^3 bacteria.

The groups of 12 mice in each case were likewise injected with 1500, 300 and 60 bacteria. Only those mice which died from day 4 to day 14 were taken into consideration in the calculation.

On the basis of the percentage of mice which survived, the strength of the vaccine was calculated using the "probit" analysis.

The test had a confidence range of 55%-192% of the reference preparation with a confidence level of 95%.

The strength of a vaccine is calculated (after "probit" transformation) using the equation

$$\frac{\text{potency, standard}}{\text{potency, test}} = \frac{\text{ED}_{50, \text{ test}}}{\text{ED}_{50, \text{ standard}}}$$

Preparation of outer membrane vesicles (OMV)

Outer membrane vesicles (OMV) were prepared by resuspending bacteria after growth and centrifuging in 10 mM Tris, pH 8.0, 2 mM EDTA. Approximately 10 ml of Tris were used to resuspend bacteria from one litre of 100 OU/ml culture.

The suspension was sonicated on ice (50% capacity, 100 W, 50% duty cycle, Branson 250 sonifier). After centrifuging for 10 min at 10,000 g, the supernatant was pelleted for one hour at 50,000 g. This pellet was resuspended in 1% (w/v) sarkosyl (sigma) in 10 mM Tris.HCl volume (Filip. C., et al., Solubilization of the cytoplasmic membrane of *Escherichia coli* by the ionic detergent sodium lauryl sarcosinate. J.Bacteriol. 255 (1973), pp. 717-722). After centrifuging for 10 min at 10,000 g, the OMVs were pelleted for one hour at 50,000 g. This method was repeated once.

The pellet finally obtained was resuspended in 10 mM Tris/HCl, pH 8.0, one-tenth of the original Tris volume. The protein concentration was determined using the BCA protein assay reagent (Pierce Chemical Company) using BSA as standard.

The desoxycholate (DOC) treatment of OMV was carried out as follows: OMVs were resuspended in 1.5% (w/v) DOC, 50 mM glycine, 5 mM EDTA, pH 9.0, sonicated for 5 min and pelleted by centrifuging: 10 min at 10,000 g followed by one hour at 100,000 g.

The LPS content of OMV is approximately 25% (wt/wt LPS/protein) and after treatment with DOC is about 10%, using KDO determination.

Characteristics of OMPs

The OMPs were characterised via trypsin sensitivity and Triton-X-100/MgCl₂ extractability as described by L. van Alphen, et al., Characteristics of major outer membrane proteins of *Haemophilus influenzae*. J.Bacteriol. 155 (1983), pp. 878-855.

Purification of antigens

- Pertussis toxin (PT) was purified from the culture supernatant using Affigelblue (Pharmacia) and column chromatography as described by Sekura, R.D., et al., Pertussis toxin, affinity, purification of a new ADP-ribosyltransferase. J.Biol.Chem. 258

(1983), 14647-14651. PT without enzymatic activity (9K, 129G) was supplied by R. Rappuoli [Pizza, M.G., et al., Mutants of pertussis toxin suitable for vaccine development, *Science* 246 (1989), 497-500].

- 5 - FHA was purified from the culture supernatant using Affigelblue and hydroxyapatite column chromatography (Sato et al., Separation and Purification of hemagglutinins from Bordetella pertussis, *Infect.Immun.* 41, (1983), pp. 313-320). Purified 69kDa OMP was supplied by R. Rappuoli (Sclavo Research Center).
- 10 - Fimbriae were isolated by means of a heat shock treatment of bacteria in the presence of urea, followed by a few centrifuging steps (Mooi, F.R., et al., Characterization of fimbriae subunits from *Bordetella* species, *Microb.Pathog.* 2 (1987) pp. 473-484).
- 15 - Endotoxin was purified by phenol extraction of bacteria (Ackers, J.L. and J.M. Dolby, The antigen of Bordetella pertussis that induces bactericidal antibody and its relationship to protection of mice, *J.Gen.Microbiol.* 70 (1972), 371-382).
- The 92kDa, 38kDa and 32kDa OMPs were obtained from the culture-cell pellet as follows:
- 20 The cells were extracted with 0.5 M CaCl_2 , 0.14 M NaCl, 1% Zwittergent 3-14 (Calbiochem) pH 4.0 (10 ml per gram wet weight of cells). After resuspension, the pH was adjusted to pH 5-6. After stirring for one hour at room temperature, the suspension was centrifuged (1 h, 3000 g)
- 25 and the supernatant was treated with 20% (v/v) ethanol, stirred for 30 minutes and centrifuged for 30 min at 10,000 g. The supernatant was dialysed against 50 mM Tris.HCl, 0.05% Zwittergent 3-14, pH 8.0. After centrifuging for 30 min at 10,000 g, the supernatant
- 30 was introduced into a DEAE Sepharose CL6B (Pharmacia) column and eluted using a linear gradient of 0-0.6 M NaCl in 50 mM Tris.HCl, 0.05% Zwittergent 3-14. Fractions enriched with either 92kDa or 32kDa were combined, concentrated with polyethylene glycol 20,000 and
- 35 precipitated with 80% ethanol. The pellets were dissolved in 50 mM Tris.HCl, 0.5% Zwittergent 3-14, pH 9.0, and passed through a Sephacryl S300 column in 50 mM Tris.

0.05% Zwittergent 3-14, pH 8.0. Fractions containing 38kDa were combined, concentrated and stored. The 38kDa OMP was purified from C-mode cells after OMV isolation. OMV pellets were treated with 5% Zwittergent 3-14, 50 mM Tris, pH 7.5, and stirred for 2 hours at 37°C. A f t e r centrifuging for 1 hour at 50,000 g, the supernatant was passed through a Sephacryl S300 column in 0.05% Zwittergent 3-14, 50 mM Tris, pH 8.0. Fractions containing 38kDa were combined, concentrated and stored.

Preparation of monoclonal antibodies

BalB/c mice were immunised i.p. with 20 µg purified antigens or OMV in the presence of 0.5 mg of AlPO_4 . (AlPO_4 gel was prepared from AlCl_3 and Na_3PO_4 in equal amounts, adjusting the pH to 7.0 using 20% (w/v) Na_2CO_3). The mice were injected four times over a period of one week. Following the final injection, the cells of the spleen were brought into contact with Sp 2/0 myeloma cells as described in Abdillahi, H and J.T. Poolman, Neisseria meningitidis group B serosubtyping using monoclonal antibodies in whole-cell ELISA. Microb.Pathog. 4, (1988), pp. 27-32 and the hybridoma cells were cultured. After adequate cell growth, the supernatants were tested via ELISA using purified antigens and OMV. Interesting hybridomas were cloned twice using the "limiting" dilution method.

Ascites fluid was obtained from these clones after introduction into the peritoneum of Pristane-injected mice. The collection of monoclonal antibodies against B.pertussis antigens is described in Poolman, J.T., et al., Description of a hybridoma bank towards Bordetella pertussis toxin and surface antigens. Microb.Pathog. 8, (1990), in press.

ELISA

One hundredth μ l of 2 μ g/ml antigen was coated in PBS (LPS in Na_2CO_3 , pH 9.0) in the wells of round-bottomed polyvinyl chloride microtitre plates. Purified OMPs were diluted to maintain the concentration of Zwittergent 3-14 below 0.001%. PT ELISA was preceded by a coating step using fetuin. The coating took place over a period of 16 hours at room temperature. The plates were washed twice with tap water + 0.02% Tween 80 (Merck). The reactions were carried out as described in Abdillahi, H. and J.T. Poolman, 1988 Neisseria meningitidis group B serosubtyping using monoclonal antibodies in whole-cell ELISA. Microb.Pathog. 4: 27-32, but 0.05% (w/v) Protifar (Nutricia) stain casein in order to block the non-specified bond. Rabbit anti-mouse peroxidase (own product) was used as second antibody.

The results were measured at 450 nm using Titertek Multiscan (Flow Labs.). "Whole-cell" ELISA was carried out as described in Abdillahi, H. and J.T. Poolman, 1988 Neisseria meningitidis group B serosubtyping using monoclonal antibodies in whole-cell ELISA, Microb.Pathog. 4: 27-32, using Protifar.

20

SDS-PAGE/immunoblotting

Purified antigens, OMV and WCV were separated by SDS polyacrylamide gel electrophoresis and labelled as described in L. van Alphen et al., 1983, Characteristics of major outer membrane proteins of Haemophilus influenzae. J.Bacteriol. 155: 878-855.

Whole cells for SDS-PAGE were obtained by suspending 3.5 ml of bacteria with 1.0 absorption at 620 nm in 70 μ g of water + 130 μ l of sample buffer. Electroblothing on nitrocellulose paper after SDS-PAGE was carried out at 140 mA constant current for a period of 1 hour on a semi-dry electroblotter (Ancos, Denmark) in accordance with the manufacturer's instructions. The paper was washed for 30 min with a physiological saline solution + 10 mM Tris, pH 7.4, containing 0.5% Tween 80 at 37°C in the same buffer. The paper was washed three times with the buffer, incubated with 0.5% Protifar in buffer for 10 min, washed and incubated with rabbit anti-mouse peroxidase for 1 hour in buffer + 0.5% Protifar. The paper was washed three times and incubated with a solution

containing 24 mg of 3,3',5,5'-tetramethylbenzidine (TMB, Sigma), 80 mg of DONS, dioctyl sulphosuccinate, dissolved in 10 ml of 96% ethanol, added to 30 ml of 10 mM Na_2HPO_4 , 5 mM citric acid, pH 5.0, to which 20 μl of 30% H_2O_2 have been added.

LEGEND

- Fig. 1: This figure shows the i.c. mouse protection value related to the 92kDa OMP/38kDa OMP ratio in the OMV vaccine.
5
- Fig. 2: OMP characterisation.
Path 1: Tohama OMV; virulent stage (X mode)
Path 2: Tohama OMV; avirulent stage (C mode)
10 Path 3: X-mode OMV after trypsin treatment
Path 4: OMV treated with Triton-X-100/MgCl₂ (supernatant)
Path 5: OMV treated with Triton-X-100/MgCl₂ (pellet)
15 Coomassie staining
- Fig. 3: OMP preparations, stained after blotting with the monoclonal mixture.
Path 1: OMV W28
20 Path 2: OMV W28 69⁻ mutant
Path 3: OMV 509
Path 4: OMV 509; C mode (avirulent stage)
Path 5: OMV 509; FHA⁻ mutant
Path 6: OMV 509; fim⁻ mutant
25 Path 7: OMV 509; stage IV
Path 8: Tohama OMV
Path 9: Tohama OMV; DOC treatment
Path 10: OMV BP-TOX6
Path 11: OMV BP-TOX6; DOC treatment
30 Path 12: OMV BP-TOX6; C mode
Path 13: OMV BP-TOX6; C mode; DOC treatment.
- Fig. 4: This figure schematically shows the (generally known) plasmid pUC18 comprising a 3.8 kb insert containing the 32 kDa OMP gene.
35

CLAIMS

1. Vaccine suitable for combating Bordetella pertussis comprising as active component, at least one or more OMPs derived from B. pertussis.
2. Vaccine, according to claim 1, in which the active component, being one or more OMPs derived from B. pertussis, are present in an OMV formulation or an artificial vesicle formulation.
3. Vaccine according to claims 1 or 2, in which the active component being one or more OMVs derived from Bordetella pertussis are present in a mixed protein-detergent formulation.
4. Vaccine according to claim 3, in which the detergent is Zwittergent 3-14.
5. Vaccine according to one or more of claims 1-4, to which an adjuvans is added.
6. Vaccine according to claim 5, in which the adjuvans is AlPO_4 .
7. Vaccine according to one or more of claims 1-6, in which the OMPs have a molecular weight of about 92kDa and 32kDa.
8. Vaccine according to claim 7, in which the OMP has a molecular weight of about 32kDa.
9. Vaccine according to claim 7 in which the OMP has a molecular weight of about 92kDa.
10. Vaccine according to claim 7, in which the OMPs have a molecular weight of 92kDa, 32kDa and 38kDa, the 92kDa OMP/38kDa OMP weight ratio being ≥ 0.25 .
11. Vaccine according to claim 10, in which the 92kDa OMP/38kDa OMP weight ratio ≥ 0.4 .
12. Vaccine according to one or more of claims 2-11, in which the OMVs have been subjected to a treatment with an effective amount of desoxycholate in order to lower the endotoxin content in the OMVs.
13. Vaccine according to one or more of claims 1-12, in which the B. pertussis strain has been chosen from the group comprising B. pertussis strain 134, B. pertussis strain 509, B. pertussis Tohama strain and B. pertussis strain BP-TOX6.

14. Vaccine according to one or more of claims 1-13, in which the OMPs are derived from Bordetella pertussis strains or other micro-organisms in which the presence of 92kDa and 32kDa OMPs has been increased via genetic engineering.
- 5 15. Vaccine according to one or more of claims 1-14, in which the OMVs are derived from fimbriae⁻ and/or FHA⁻ B.pertussis strains.
16. Vaccine according to claim 15, in which the OMVs are derived from the B.pertussis strain BP 509, fim⁻ mutant or the
- 10 B.pertussis strain BP 509, FHA⁻ mutant.
17. Vaccine according to one or more of claims 1-16, in which PT toxoid or FHA, or PT toxoid and FHA, are present, optionally in the form of a divalent conjugate, as additional active component(s).
- 15 18. Vaccine according to claim 17, in which the PT toxoid used in the vaccine has been obtained by means of a formaldehyde treatment or glutaraldehyde treatment of PT.
19. Vaccine according to claim 17, in which the PT toxoid has been derived from a B.pertussis strain in which the PT gene has
- 20 been modified in such a way that it codes for an immunologically active PT having no toxic effect or a toxic effect acceptable for vaccine use.
20. Vaccine according to claim 17, in which the FHA has been purified with the aid of chromatography using hydroxyapatite.
- 25 21. Method for the vaccination of humans, in particular children less than 2 years old, in which an effective amount of the vaccine according to one or more of Claims 1-20 is used.
22. Method according to claim 21, in which the amount of vaccine to be injected contains a dose of 5-25 µg OMV.
- 30 23. OMPs and OMVs derived from B.pertussis, suitable for use in the preparation of vaccines, defined in one or more of Claims 1-20.
24. OMVs according to Claim 23, the OMVs having been subjected to a treatment with an effective amount of desoxycholate
- 35 in order to lower the endotoxin content in the OMVs.

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

NL 9100185
SA 51997

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 30/01/92. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family members)	Publication date
WO-A- 9012086	18-10-90	AU-A- 5432690	05-11-90
		CA-A- 2013571	30-09-90
		CN-A- 1046532	31-10-90

WO-A- 9115505	17-10-91	None	

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATION WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹

This International search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim numbers _____ because they relate to subject matter not required to be searched by this Authority, namely: _____
 Although claims 21 and 22 are directed to a method of treatment of the human/animal body the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claim numbers _____ because they relate to parts of the International application that do not comply with the prescribed requirements to such an extent that no meaningful International search can be carried out, specifically _____
3. ☐ Claim numbers _____ because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ²

This International Searching Authority found multiple inventions in this International application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International search report covers all searchable claims of the International application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this International search report covers only those claims of the International application for which fees were paid, specifically claims: _____
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers: _____
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	Journal of Bacteriology, volume 166, no. 1, April 1986, American Society for Microbiology; S.K. Armstrong et al.: "Bordetella pertussis major outer membrane porin protein forms small, anion-selective channels in lipid bilayer membranes", pages 212-216, see page 212, right-hand column, line "Bacteria and culture conditions..." to page 213, left-hand column, line 39 ---	23-24
X	Reviews of Infectious Diseases, volume 10, supplement 2, July-August 1988, The University of Chicago; C.D. Parker et al.: "Surface proteins of Bordetella pertussis", pages S327-S330, see page S327, right-hand column, line "Isolation and analysis of cell...." to page S328, right-hand column, last line ---	23-24
X	Proceedings of the National Academy of the USA, volume 86, no. 10, May 1989, (Washington, DC, US) I.G. Charles et al.: "Molecular cloning and characterization of protective outer membrane protein P.69 from Bordetella pertussis", pages 3554-3558, see page 3554, right-hand column, first paragraph ---	1,23,24
E	WO,A,9115505 (CONNAUGHT LABORATORIES LIMITED) 17 October 1991, see page 12, example 9 -----	1,5,6, 17,23

INTERNATIONAL SEARCH REPORT

International Application No


PCT/NL 91/00185

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁸		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl.5 C 07 K 13/00 A 61 K 39/10		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl.5	A 61 K C 07 K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X, P	WO, A, 9012086 (WASHINGTON UNIVERSITY) 18 October 1990, see page 6, line 2 - page 9, line 12 ---	1
X	--- Biological Abstracts, volume 88, 1989, M.G. Thomas et al.: "Human serum antibody responses to Bordetella pertussis infection and pertussis vaccination", abstract no. 4699, & J. Infect. Dis 159(2): 211-218, 1989, see the abstract ---	23-24
X	--- Journal of Bacteriology, volume 159, no. 2, August 1984, American Society for Microbiology; F. Shareck et al.: "Cloning of Bordetella pertussis outer membrane proteins in Escherichia coli", pages 780-782, see the whole article --- -/-	23-24
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
09-12-1991		06.02.92
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		Nicole De Bie

I.C.POTENCY <---> 92K/38K RATIO


fig-1

+ 92K/38 ratio




THIS PAGE BLANK (USPTO)

fig - 2




THIS PAGE BLANK (USPTO)

fig-3



THIS PAGE BLANK (USPTO)

fig-4



THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)